


Space Weather

Lecture 9: Ionosphere



Elena Kronberg (room 442)
elena.kronberg@lmu.de


Some history



- In 1839, Carl Friedrich Gauss speculated that an electrically conducting region of the atmosphere could account for observed variations of Earth's magnetic field.
- The name “ionosphere” was formally defined in 1950 by a committee of the Institute of Radio Engineers as “the part of the Earth's upper atmosphere where ions and electrons are present in quantities sufficient to affect the propagation of radio waves.”

Ionosphere formation



- Ionosphere is the transition region between fully ionized magnetospheric plasma and neutral atmosphere.
- It is a mixture of ionized and neutral particles: partially ionized plasma.
- Therefore, Coulomb and neutral collisions may contribute to electrical conductivity.



- Fully ionized plasmas: magnetosphere, solar wind; partially ionized plasma: Sun's photosphere, chromosphere, ionosphere
- There are two main sources of ionization: ultraviolet radiation from the Sun and precipitation of energetic particles from the magnetosphere into the atmosphere.

Solar Ultraviolet (UV) Ionization


- The photoionization layer in the ionosphere exhibits a strong dependence on geographic latitude, time of day and season.



Credit: Baumjohann+Treumann

Ionization by Energetic Particles

- Dominates at high magnetic latitudes in the auroral zone, where photoionization becomes less important
- During nighttime, when photoionization ceases, ionization due to particle impact can maintain the ionosphere




Credit: Baumjohann+Treumann

Recombination and Attachment


- The production of ionization in the ionosphere either by solar UV radiation or by energetic particles would, if it continued endlessly, lead to full ionization of the upper atmosphere.
- In reality two processes counteract the ionization:
 - 1 Recombination of ions and electrons to reform neutral atoms
 - 2 Attachment of electrons at neutral atoms or molecules to form negative ions

Penetration of electro-magnetic radiation and altitude



Credit: STScI/JHU/NASA

Ionospheric layers: from ~ 60 to ~ 1000 km




- D-layer is connected with the most energetic precipitation.
- E-layer contains mainly O_2^+ and NO^+ ions produced by UV radiation (100–150 nm) and solar X-rays (1–10 nm).
- F₁-layer is composed of mainly O^+ produced by UV radiation in the range from about 17 to 91 nm.
- F₂-layer is also composed of O^+ , formation is complicated: photochemical processes, vertical movement due to neutral drag and magnetospheric effects.

F-layer is the most important region for long distance HF radio communications.

Credit: Paschmann+02

Ionospheric layers and HF radio communication





© 1999 Encyclopædia Britannica, Inc.

Space weather effects: Associated with solar flare

A Space Weather event: Sept 6, 2017

- The sudden outburst of electromagnetic energy travels at the speed of light. The sunlit side of Earth is exposed. The increased level of X-ray and extreme ultraviolet (EUV) radiation results in ionization.
- The D-layer becomes more dense. This can cause HF radio signals to become degraded or completely absorbed.



During a mild geomagnetic storm on Feb 4, 2022 40 from 49 Starlink satellites were burned in the atmosphere

- Ionospheric dynamics was not well predicted.
- Atmospheric drag is caused by frequent collisions of molecules with a satellite. It reduces the altitude of satellite's orbit and leads to re-entry.





Source: ROOM

Ionospheric conductivity

- The presence of free charges in the ionosphere results in a conductivity which is anisotropic due to the effects of the magnetic field and collisions.
- The three current components are:


- ① the field-aligned (parallel) current $j_{\parallel} = \sigma_{\parallel} E_{\parallel}$
- ② the Pedersen current $j_P = \sigma_P E_{\perp}$ flowing parallel to the transverse electric field E_{\perp}
- ③ the Hall current $j_H = \sigma_H \hat{b} \times E$ which flows transverse to both the magnetic and electric fields



Credit: Menk&Waters + Paschmann+02

Ionospheric conductivity


The conductivities are defined as $\sigma_{||} = \sum_s \frac{n_s q_s^2}{m_s \nu_s}$, $\sigma_P = \sum_s \frac{n_s q_s^2}{m_s} \frac{\nu_s}{\nu_s^2 + \omega_{cs}^2}$ and $\sigma_H = - \sum_s \frac{n_s q_s^2}{m_s} \frac{\omega_{cs}}{\nu_s^2 + \omega_{cs}^2}$, where n_s , q_s , m_s and ν_s indicate density, charge, mass and particle-neutral collision frequency, respectively, of particles of species s , and $\omega_{cs} = q_s B / m$ is the gyrofrequency of species s .



Credit: Paschmann+02



Plasma Convection

- Flux tube and plasma convection caused by magnetic merging




Credit: Baumjohann+Treumann

Reflection of the convection in the ionosphere



Credit: Paschmann+02, Kivelson&Russell


Currents in the ionosphere



Credit: Baumjohann+Treumann


Currents in the ionosphere

AMPERE (Active Magnetosphere and Planetary Electrodynamics Response Experiment) observations



Credit: JHU/APL


Ionosphere-Magnetosphere coupling



Le+2010; also Wikipedia

Currents during substorm


- Left side: convection electric field during substorm strengthens convection auroral electrojets (Hall currents, travel around the E region)
- Right side: substorm electrojet is formed by the substorm current wedge (Hall current)
- The strength of substorm electrojet is mainly determined by increase in ionospheric conductance because of strong particle precipitation.



Credit:
Cravens, 1997


Substorm Current Wedge

- The substorm current wedge diverts part of the neutral sheet current along magnetic field lines through the ionosphere




McPherron&Chu 16

Aurora: colors



Collision of particles at different energies with ionospheric atoms lead to emissions of different colors





Aurora: forms

Copyright: Off the map travel



Chromosphere and Ionosphere: analogy



Particle precipitation triggers ionization and release of energy → Solar flare

Credit: SSL Berkely

Summary

- The ionosphere is strongly affected by the dynamics of the Sun and particle precipitation in the atmosphere.
- The magnetosphere and ionosphere are coupled through the motion of particles along field lines producing field-aligned currents.
- These currents characterize the location of auroras.
- Intense Hall currents result in auroral electrojets that flow toward midnight around the auroral ovals (lead to irregular variations of the ground magnetic field on scales of seconds to days).
- These currents lead to geomagnetically induced currents which may disturb, e.g., work of power grids...

Literature

- W. Baumjohann and R. Treumann, Basic Space Plasma Physics, 1996
- M. Kivelson and C. Russell, Introduction to Space Physics, 1995
- A. Brekke, Physics of the Upper Polar Atmosphere, 2013
- G. Paschmann, S. Haaland and R. Treumann, Auroral Plasma Physics, 2002
- F. Menk and C. Waters, Magnetoseismology: Ground-based remote sensing of Earth's magnetosphere, 2013